

Effect of Some Bio-Regulators on Growth and Yield of Some Wheat Varieties under Newly Cultivated Land Conditions

Ali A El-Hosary¹, Gaber Y Hammam¹, Abdalla El-Morsi², Esmat A Hassan², Mohamed E El-Awadi^{2,*},
Yasser R Abdel-Baky²

¹ Agronomy Department, Faculty of Agriculture, Benha University, Egypt

² Botany Department, National Research center, Egypt.

ABSTRACT

This investigation was conducted to study the effect of some bio-regulators, tryptophan, cysteine, thiamine, ascorbic acid and yeast extract on growth, yield and its components and some chemical constituents of some wheat varieties; Sakha93, Gemiza7 and Gemiza9 under newly cultivated land conditions. For this purpose two field experiments were carried out during two successive growing seasons (2008/2009 and 2009/2010) at the experimental farm of National Research Center at Nubaria, Egypt. Results revealed that foliar application of different applied growth bio-regulating substances, with different concentrations, significantly increased all growth criteria, plant height, number of tillers and leaves per plant, fresh and dry weight of plant. Grain and straw yield of wheat plants and yield components were also significantly increased as result of foliar application of the used bioregulators. On the other hand Gemiza-7 variety was most effective of all characters under study compared with the other two varieties Gemiza-9 and Sakha-93.

Key Words

Wheat, Bio-regulators, Growth, Pigments, Yield, Yield components.

Correspondence to:

Mohamed E El-Awadi
Botany Department,
National Research center,
Egypt.
E-mail: el-awadi@yahoo.com

1 Introduction

Wheat is one of the major cereal crops with a unique protein, which is consumed by humans and is grown around the world in diverse environments (Mac Ritchie, 1994). Wheat provides approximately one fifth of the calories in the human diet and is an important source of vegetable protein and nutrients for a large proportion of the world's population. Modern wheat varieties differ in their grain concentrations of N, Zn and Fe. (Cakmak et al., 2000). (Sultan et al., 2000) reported, Wheat cultivars differed in yield and its components. Also Zaki, et al., (2004) established that, grain, straw and biological yields and its components were significantly differed owing to varietal differences. In addition Sharaan, (2000) found that wheat varieties were significantly different in plant height, spike length, number of spikes, number of grains per spike and grain weight per spike Zaki et al., (2007) reported that, there were significant differences in all growth characters of wheat cultivars, as follows i.e. (Sakha93, Gemiza7 and Gemiza9) under their study.

Plant growth regulators (PGRs) are actively involved in a multitude of metabolic processes and play essential roles in plant growth and development under both stress and natural conditions. They also act as chemical messengers to modulate various processes or genes involved in plant growth and development (Morgan, 1990).

Plant growth regulators (PGRs) also

play important roles in plant adaptation to stressful environments, including drought stress (Huang et al., 2008).

Under both wild and cultivated conditions, plants often experience a multitude of environmental stresses such as drought, salinity, waterlogging, extremes of temperature, and mineral toxicities and deficiencies. Environmental stresses are undoubtedly a major cause of food insecurity in many countries around the world, in particular, in developing countries where there is a major challenge to produce sufficient food. A large proportion of the world, agriculture depends on rainfall for irrigation, as good quality water supply is highly limited or unpredictable. In many of such regions, crops are often negatively affected by severe drought. It has been estimated that the countries that generate two third of the world's agricultural product experience water-deficit conditions on a regular basis (Revenge et al., 2000).

In plants, amino acids fulfill a wide variety of functions. Their common role is to serve as building blocks of proteins, which exert manifold functions in plant metabolism, and as metabolites and precursors involved in plant defense, vitamin, nucleotide and hormone biosynthesis, and as precursors of a huge variety of secondary metabolites. One way or the other, as active catalysts or as precursors, amino acids are essentially involved in all metabolic, regulatory, and physiological aspects of plant metabolism (Buchanan et al., 2000).

Unlike humans, plants are capable of producing vitamins. In fact, the foods

highest in vitamins are plant-based, such as raw fruits and vegetables. Vitamins could be considered as bio-regulators compounds which in low concentrations exerted a profound influence upon plant growth. In general, the energy metabolic pathway could be affected by one or another of these substances. Vitamins could serve as coenzyme in decarboxylation of α -keto acids, such as pyruvic acid and keto-glutamic acid which has its importance in the metabolism of carbohydrates and fats (Bidwell, 1979).

Vitamins are important cofactors for the transketolation reactions of the pentose phosphate cycle, which provides pentose phosphate for nucleotide synthesis and for the reduced NADP required in various synthetic pathways (Kawasaki, 1992).

On the other hand, Amer, (2004) and Kurtzman and Fell (2005) demonstrated that, Active dry yeast is a natural safety biofertilizers causes various promotive effect on plants. It is considered as a natural source of cytokinins which simulates cell division and enlargement as well as the synthesis of protein, nucleic acid and vitamin-B. It also releases CO₂ which reflected in improving net photosynthesis.

The objectives of this study was to evaluate the beneficial effect of some bio-regulators as foliar application on growth, yield and yield components and some chemical constituents of some wheat varieties grown under newly reclaimed soil conditions.

2 Materials and Methods

This investigation was carried out at the experimental farm of National Research Center at Nubaria, Egypt, during two successive seasons 2008\2009 and 2009\2010, to study the effect of some bio-regulators on growth development, yield and yield components of three cultivars of wheat plant. (*Triticum aestivum* L.).

Plant Materials

Pure lines and uniform grains of three local cultivars of wheat plant (*Triticum aestivum* L.) Sakha93,

Gemiza-7 and Gemiza-9. These cultivars were obtained from Wheat Research Department, Agricultural Research Center, Ministry of Agriculture, Egypt.

Chemical materials

Three types of chemical materials were used, tryptophan (C₆H₄NH.CH:C) and cysteine (C₃H₇NO₂S) as amino acids, ascorbic acid (Vitamin-C) (C₆H₈O₆) and thiamine (Vitamin-B1) as vitamins and baking yeast (*Saccharomyces cerevisiae*) extract.

The experiments were carried out under sandy soil conditions. Grains of the three cultivars were sown at 27 and 23 November for the first and the second season respectively. The experimental design was split plot design with three replications. Each experiment included 33 plots and the plot area was 10.5 m² (1/400 of feddan), three meters in length and three and half meters in width. Each plot contained twenty rows. Seeds were planted at the rate of 60 Kg/feddan using drilling and cross rows method, the spacing between the rows was 15cm. The treatments of the bio-regulators were distributed at random in the main plot, and the three varieties were assigned in the sub plot randomly too.

Nitrogen fertilizer was applied at the rate of 120 Kg N/feddan in the form of ammonium nitrate (33.5%N) and was divided in ten equal portions. The dosage was added before the irrigation.

Phosphorus fertilizer was applied before sowing (during land preparation) at rate of 50 Kg P₂O₅/feddan in the form of calcium super phosphate (15.5% P₂O₅).

Potassium sulphate was added at rate of 50 Kg per feddan, and applied one month after sowing.

Two concentrations from each bio-regulator (50 and 100 mg l⁻¹ of tryptophan, thiamine and ascorbic acid) and 100 and 150 mg l⁻¹ of cysteine as well as 1 and 2 g l⁻¹ of yeast extract were added twice to the plants as foliar spray in addition to control plants which sprayed with distilled water. First spray was carried out at vegetative stage (30 days after sowing) the spray volume of this sprinkling was 200 liter per feddan, and the second spray was seven days later,

its volume was 250 liters per feddan.

The sample was taken fifteen days after the second spray (at heading stage).

Growth and yield measurements

- Plant height (cm).
- Number of tillers per plant.
- Number of leaves per plant.
- Fresh weight of shoot per plant.
- Dry weight of shoot per plant.
- Flag leaf area (cm²)

Yield and yield components characters

At harvest, ten plants from each plot were chosen randomly to estimate the following data.

- Spike length (cm).
- Number of spikes per plant.
- Spike weight (g).
- Number of grains per spike.
- Grain weight per spike (g).
- Grain yield per feddan (Kg) which

was determined from the whole grain yield of each sub-plot and converted to tons of grains per feddan.

Photosynthetic pigments

Chlorophylls a, b and carotenoids were extracted from fresh leaf tissues and estimated, colorimetrically, according to method described by Wettstein (1957).

Statistical Analysis

Each experiment was statistically analyzed as a split plot design according to Snedecor and Cochran, (1980). Combined analysis of the two seasons was carried out whenever homogeneity of variance was detected for all studied traits. The Duncan multiple range test was used to compare the treatment means (Duncan, 1955). The MSTATC (1989) program was used in this connection.

3 Results

Growth characters

Effect of bio-regulators

According to data in Table (1), yeast extract at concentration of 1000 mg l⁻¹ was the most effective treatment increasing wheat plant height followed by Cysteine treatment at concentration of 150 mg l⁻¹. Data also show that there was no significant difference between the treatments 150 mg l⁻¹ of cysteine and 1000 mg l⁻¹ of yeast extract in the second season.

The highest number of tillers and leaves per plant was obtained by using tryptophan at concentration of 50 mg l⁻¹. Data show also that, the highest values of fresh weight of plant were obtained by treatment of 1000 mg l⁻¹ yeast in the second season and combined data of the two seasons. On the other hand treatment of 100 mg l⁻¹ ascorbic acid recorded the highest value in the first season whereas, the untreated plants (control) recorded the lowest values of the fresh weight of plant during first, second season and their combined data.

It could be emphasized from the present data that, dry weight of wheat plant was significantly affected by all bio-regulating substances at different concentrations in both season and their

combined data. Results show that, the effect was strikingly noticeable with the treatment of ascorbic acid 100 mg l⁻¹ in the first season and tryptophan 50 mg l⁻¹ in the second season and the combined data.

It is clear from Table (2) that, flag leaf area was significantly increased as a result of foliar application with the used bio-regulating substances at different concentrations. Ascorbic acid treatment at 100 mg l⁻¹ recorded the highest value in this respect, while, the lowest value obtained by untreated plants (control).

Varieties performance

Data in Tables (1 and 2) revealed that, the examined varieties significantly varied in their responses to foliar spray with the different bio-regulating

substances. Gemiza-7 variety gave the highest values of plant height, fresh and dry weight of plant and flag leaf area. While, Gemiza-9 variety recorded the highest values of number of tiller and leaves per plant and these results were true for the two seasons and their combined data.

Effect of interaction

The results in Tables (3 and 4) cleared that, there were significant increases of all growth characters under study in the two seasons and their combined data as affected by interaction between wheat varieties and bio-regulating substances.

The highest values of plant height were obtained from the treatment 1000 mg l⁻¹ of yeast with Gemiza-7 variety, during the first, second season

Table 1. Plant height, number of tillers and number of leaves per plant, as affected by some growth bio-regulators treatments and varieties of wheat at heading stage during two growing seasons and their combined data.

Treatments	Plant height (cm)			No. of Tillers/plant			No. of leaves/plant			
	S1	S2	Comb.	S1	S2	Comb.	S1	S2	Comb.	
Control	55.97 d	58.40 h	57.19 e	2.334 e	3.008 f	2.671 f	11.17 g	12.33 g	11.75 h	
Tryptophan	50 mg l ⁻¹	74.70 ab	77.81 c-e	76.26 c	6.017 a	7.563 a	6.790 a	23.14 a	26.35 a	24.74 a
	100 mg l ⁻¹	71.60 c	74.41 g	73.00 d	4.533 c	5.158 d	4.846 d	17.33 ef	18.90 f	18.11 g
Cystine	100 mg l ⁻¹	70.74 c	76.43 f	73.58 d	3.691 d	4.250 e	3.971 e	18.03 de	20.78 de	19.40 f
	150 mg l ⁻¹	75.73 ab	80.67 a	78.20 ab	5.034 bc	5.750 cd	5.392 c	21.49 b	22.43 c	21.96 c
Thiamine	50 mg l ⁻¹	73.80 b		76.34 c	5.046 bc	5.456 cd	5.251 cd	19.75 c	20.58 e	20.17 e
	100 mg l ⁻¹	69.51 c	77.09 d-f	73.30 d	4.670 bc	5.264 cd	4.967 cd	18.43 d	20.21 e	19.32 f
Ascorbic Acid	50 mg l ⁻¹	75.10 ab	79.47 b	77.28 bc	4.612 bc	5.482 cd	5.047 cd	18.20 d	20.15 e	19.17 f
	100 mg l ⁻¹	71.28 c	76.80 ef	74.04 d	5.209 b	6.509 b	5.859 b	20.09 c	21.74 cd	20.91 d
Yeast	1000 mg l ⁻¹	76.26 a	80.77 a	78.51 a	5.123 bc	6.611 b	5.867 b	21.10 b	24.98 b	23.04 b
	1500 mg l ⁻¹	69.99 c	78.27 b-d	74.13 d	4.867 bc	5.904 c	5.386 c	16.91 f	22.29 c	19.60 ef
Sakha-93		±1.31 cm	66.74 c	66.50 c	4.706 ab	5.642 a	5.174 a	17.72 c	19.97 c	18.84 c
Gemiza-7		±8.65 cm	85.39 a	81.16 a	4.432 b	5.275 b	4.853 b	18.65 b	20.50 b	19.58 b
Gemiza-9		±1.31 cm	76.68 b	73.75 b	4.808 a	5.708 a	5.258 a	19.72 a	22.46 a	21.09 a

S1 = first season, S2 = second season, Comb. = combined data.

Table 2. Fresh and dry weight per plant and flag leaf area, as affected by some growth bio-regulators treatments and varieties of wheat at heading stage during two growing seasons and their combined data.

Treatments	Fresh weight/plant (g)			Dry weight/plant (g)			Flag leaf area (cm ²) (Average two seasons)	
	S1	S2	Comb.	S1	S2	Comb.		
Control	10.53 f	14.00 h	12.27 h	8.797 f	9.558 g	9.178 g	26.20 g	
Tryptophan	50 mg l ⁻¹	27.20 d	38.78 c	32.99 bc	17.76 b	23.00 a	20.38 a	41.60 b
	100 mg l ⁻¹	22.58 e	31.42 e	27.00 f	12.24 e	15.28 ef	13.76 f	37.75 e
Cystine	100 mg l ⁻¹	21.87 e	28.14 g	25.00 g	15.06 d	16.02 de	15.54 de	37.34 e
	150 mg l ⁻¹	30.04 b	35.59 d	32.81 bc	16.36 c	21.04 b	18.70 bc	40.19 c
Thiamine	50 mg l ⁻¹	29.27 bc	34.91 d	32.09 c	14.52 d	16.75 d	15.64 de	38.94 d
	100 mg l ⁻¹	28.76 bc	29.75 f	29.26 e	14.68 d	15.85 d-f	15.26 de	41.67 b
Ascorbic Acid	50 mg l ⁻¹	28.12 cd	32.63 e	30.37 d	16.99 bc	14.74 f	15.87 d	39.92 c
	100 mg l ⁻¹	33.67 a	40.77 b	37.22 a	19.39 a	16.84 d	18.12 c	44.67 a
Yeast	1000 mg l ⁻¹	32.42 a	43.62 a	38.02 a	19.01 a	19.81 c	19.41 b	38.14 e
	2000 mg l ⁻¹	28.17 cd	38.64 c	33.41 b	14.50 d	15.51 ef	15.01 e	36.53 f
Sakha-93		23.03 c	27.83 c	25.43 c	14.07 c	5.642 a	13.72 c	33.11 c
Gemiza-7		29.27 a	38.39 a	33.83 a	19.09 a	5.275 b	18.12 a	45.73 a
Gemiza-9		27.51 b	34.21 b	30.86 b	17.13 b	5.708 a	16.40 b	36.52 b

S1 = first season, S2 = second season, Comb. = combined data.

and their combined data.

Treatment of 50 mg^l⁻¹ tryptophan with Gemiza-9 variety recorded the highest values of number of tillers in the first, second season and their combined data. While, 50 mg^l⁻¹ treatment gave the highest number of leaves per plant when used on Sakha-93 variety during the same periods. On the other hand, treatment of 150 mg^l⁻¹ cysteine with Gemiza-7 variety recorded the highest value of fresh weight of plant in the first season. While, ascorbic acid at 100 mg^l⁻¹ recorded the highest values of this character when used on Gemiza-9 variety during second season and combined data.

The treatment of 50 mg^l⁻¹ tryptophan was more effective in increasing dry weight of plant during the second season and the combined data. While ascorbic acid treatment at 100 mg^l⁻¹ was more effective in the first season when used on Gemiza-9 variety. The most effective

treatments for increasing flag leaf area was 150 mg^l⁻¹ cysteine when used on Gemiza-7 variety followed by thiamine treatment at 100 mg^l⁻¹ on Gemiza-7 variety also which was no significant difference between them.

Yield and yield components

Effect of bio-regulators

Foliar application of all concentrations of bioregulating substances under study (Tables 5-6) resulted in significant increases in spike length, number of spike per plant, weight of spike, number of grains per spike, number of spikelet per spike, number of grains per spikelet, grains weight per spike, grain and straw yield of wheat plant, during the two growing seasons and their combined data compared with untreated plants (control).

Tryptophan treatment at 50 mg^l⁻¹ recorded the highest values of spike length in the combined data. While, concentration of 100 mg^l⁻¹ was

the best treatment in the first season. On the other hand 100 mg^l⁻¹ thiamine treatment recorded the highest value in this character during second season. Data also show that, yeast extract at 1000 mg^l⁻¹ concentration gave the highest values of number of spike per plant during first and combined data while, the highest value was obtained by 50 mg^l⁻¹ tryptophan treatment in the second season.

Also, tryptophan treatment at 50 mg^l⁻¹ resulted in the highest values of weight of spike per plant and grains weight per spike during the two growing seasons and their combined data as well as grain yield and number of grains per spike during the first season and the combined data, whereas 1000 mg^l⁻¹ yeast application recorded the highest value in the second season. In addition there was no significant difference between the two treatments in the second season.

Straw yield was positively affected

Table 3. Plant height, number of tillers and number of leaves per plant, as affected by interaction between some growth bio-regulators treatments and varieties of wheat at vegetative growth stage during two growing seasons and their combined data.

Treatments		Plant height (cm)			No. of Tillers/plant			No. of leaves/plant			
		S1	S2	Comb.	S1	S2	Comb.	S1	S2	Comb.	
Control	Sakha-93	37.1 n	35.5 q	36.3 o	1.07 l	1.60 kl	1.33 n	5.32 no	5.83 mn	5.58 o	
	Gemiza-7	42.7 m	36.8 p	39.7 n	1.33 l	1.20 l	1.27 n	5.15 o	5.34 n	5.24 o	
	Gemiza-9	37.5 n	41.3 o	39.4 n	1.75 l	2.40 jk	2.07 m	6.68 n	6.90 m	6.79 n	
Tryptophan	50 mg ^l ⁻¹	Sakha-93	53.9 c-e	53.3 d	53.6 c	4.47 ab	5.15 ab	4.81 ab	15.92 b-d	18.85 b	17.38 bc
		Gemiza-7	57.4 ab	56.9 a	57.2 a	3.53 c-h	4.33 b-e	3.93 c-g	11.12 i-l	11.40 jk	11.26 jk
		Gemiza-9	48.9 f-i	50.9 f	49.9 ef	3.97 ad	5.00 ab	4.48 a-c	12.26 hi	15.82 ef	14.04 fg
	100 mg ^l ⁻¹	Sakha-93	42.9 m	42.5 n	42.7 m	2.60 ik	3.03 g-j	2.81 j-l	10.51 j-m	11.24 j-l	10.88 j-l
		Gemiza-7	49.1 f-h	52.8 d	50.9 e	2.80 h-k	3.57 e-i	3.18 h-l	9.877 lm	10.67 j-l	10.27 lm
		Gemiza-9	45.1 l	44.4 lm	44.8 l	2.93 g-k	4.16 b-g	3.55 d-i	11.06 i-l	11.87 j	11.47 j
Cysteine	50 mg ^l ⁻¹	Sakha-93	49.7 fg	46.5 jk	48.1 g-i	2.57 jk	2.68 ij	2.62 lm	9.253 m	10.07 l	9.66 m
		Gemiza-7	54.1 cd	55.9 a-c	55.0 b	2.47 k	3.49 e-j	2.98 i-l	11.17 i-l	13.80 hi	12.49 hi
		Gemiza-9	46.3 j-l	44.8 lm	45.6 kl	2.90 g-k	3.87 c-h	3.38 f-k	12.09 hi	14.47 gh	13.28 gh
	100 mg ^l ⁻¹	Sakha-93	52.0 e	49.3 gh	50.7 e	3.80 a-f	4.10 b-g	3.95 c-g	12.59 g-i	15.54 fg	14.06 fg
		Gemiza-7	57.8 a	56.6 ab	57.2 a	3.47 d-h	4.04 b-g	3.75 d-h	14.50 d-f	16.27 ef	15.38 e
		Gemiza-9	48.5 f-i	45.5 kl	47.0 ij	3.87 a-e	4.56 a-e	4.21 b-d	19.77 a	20.24 a	20.01 a
Thiamine	50 mg ^l ⁻¹	Sakha-93	48.6 f-i	52.1 de	50.3 e	4.30 a-c	4.80 a-d	4.55 a-c	15.21 c-e	17.70 cd	16.45 d
		Gemiza-7	55.8 bc	52.9 d	54.3 bc	3.40 d-i	3.20 f-j	3.30 g-l	11.47 i-k	11.30 j-l	11.38 jk
		Gemiza-9	49.9 f	48.3 hi	49.1 fg	3.60 c-h	3.86 c-h	3.73 d-h	14.80 de	16.39 ef	15.60 e
	100 mg ^l ⁻¹	Sakha-93	46.0 j-l	50.3 fg	48.1 g-i	3.00 f-k	3.17 f-j	3.08 h-l	10.23 j-m	11.78 j	11.01 j-l
		Gemiza-7	53.3 de	51.3 ef	52.3 d	2.87 g-k	2.80 h-j	2.83 j-l	9.77 lm	10.35 kl	10.06 lm
		Gemiza-9	47.3 h-k	47.3 ij	47.3 hi	3.68 b-g	3.60 e-i	3.64 d-i	13.97 e-g	15.90 ef	14.93 ef
Ascorbic Acid	100 mg ^l ⁻¹	Sakha-93	46.9 i-l	49.4 gh	48.1 g-i	3.97 a-d	4.07 b-g	4.02 c-f	12.47 hi	14.40 hi	13.43 gh
		Gemiza-7	54.4 cd	54.9 c	54.7 bc	4.00 a-d	4.37 b-e	4.18 b-e	13.13 f-h	16.80 de	14.9 ef
		Gemiza-9	100 mg ^l ⁻¹	44.1 m	44.7 l	3.43 d-h	3.60 e-i	3.52 e-i	11.77 h-j	13.41 hi	12.59 h
	100 mg ^l ⁻¹	Sakha-93	45.3 kl	48.0 i	47.2 i	4.30 a-c	4.90 a-c	4.60 a-c	16.42 bc	19.10 b	17.76 b
		Gemiza-7	58.6 a	56.4 ab	57.5 a	4.10 a-d	4.80 a-d	4.45 a-c	15.32 c-e	18.39 bc	16.85 cd
		Gemiza-9	47.7 g-j	50.1 fg	48.9 fg	4.58 a	5.50 a	5.04 a	19.70 a	20.36 a	20.03 a
Yeast	1000 mg ^l ⁻¹	Sakha-93	47.2 h-k	49.5 gh	48.4 gh	3.13 e-k	3.70 d-i	3.42 f-j	10.10 k-m	13.20 i	11.65 ij
		Gemiza-7	52.6 de	56.8 a	54.7 bc	4.00 a-d	4.23 b-f	4.11 c-e	16.92 b	19.30 ab	18.11 g
		Gemiza-9	45.4 kl	46.6 jk	45.9 jk	3.33 d-j	3.84 c-h	3.59 d-i	14.54 d-f	15.96 ef	15.25 e
	2000 mg ^l ⁻¹	Sakha-93	47.1 h-l	150 mg ^l ⁻¹	45.9 jk	2.53 jk	2.90 h-j	2.72 kl	9.60 lm	11.01 j-l	10.31 lm
		Gemiza-7	53.7 de	100 mg ^l ⁻¹	54.6 bc	3.60 c-h	2.87 h-j	3.23 h-l	9.86 lm	11.03 j-l	10.45 k-m
		Gemiza-9	45.1 l	74.41 g	45.3 kl	3.13 e-k	3.04 g-j	3.09 h-l	12.31 hi	14.37 g-i	13.34 gh

S1 = first season, S2 = second season, Comb. = combined data.

by 100 mg^l⁻¹ ascorbic acid treatment during the first, second season and their combined data compared with the untreated plants (control).

Varieties performance

The examined cultivars significantly

varied in their response to different bio-regulators in the yield and yield component characters (Tables 5-6). In comparison between varieties, Gemiza-7 variety significantly exceeded in spike length, weight of spike in the two

growing seasons and their combined data, number of grains per spike in the second season, grain weight of spike and grain yield in the two growing season and their combined data.

Gemiza-9 variety recorded the

Table 4. Fresh and dry weight of plant, as affected by interaction between some growth bio-regulators treatments and varieties of wheat at vegetative growth stage during two growing seasons and their combined data.

Treatments		Fresh weight/plant (g)			Dry weight/plant (g)			
		S1	S2	Comb.	S1	S2	Comb.	
Control	Sakha-93	5.71 q	6.770 s	6.240 o	0.973 q	1.577 n	1.275 o	
	Gemiza-7	6.62 pq	7.467 s	7.043 o	1.160 pq	1.727 mn	1.443 o	
	Gemiza-9	6.95 pq	9.150 r	8.052 n	1.490 op	2.088 lm	1.789 n	
Tryptophan	50 mg ^l ⁻¹	Sakha-93	17.42 b	20.97 b	19.20 b	3.777 bc	4.467 bc	4.122 bc
		Gemiza-7	15.92 bc	19.62 c	17.77 d	3.530 cd	4.357 bc	3.943 c
		Gemiza-9	15.27 cd	16.10 g-i	15.68 e-g	3.173 de	3.207 f-i	3.190 d
	100 mg ^l ⁻¹	Sakha-93	10.05 k-n	10.66 pq	10.35 l	1.950 j-n	2.083 lm	2.017 l-n
		Gemiza-7	11.84 g-k	12.62 mn	12.23 jk	2.083 j-m	2.717 jk	2.400 i-k
		Gemiza-9	11.32 h-k	12.12 no	11.72 k	2.363 h-k	2.750 i-k	2.557 h-j
Cysteine	50 mg ^l ⁻¹	Sakha-93	8.680 no	9.390 r	9.035 m	1.863 l-o	1.957 l-n	1.910 mn
		Gemiza-7	13.98 d-f	15.49 hj	14.73 gh	2.380 h-j	3.533 ef	2.957 d-f
		Gemiza-9	10.24 j-n	10.47 q	10.35 l	1.627 no	2.687 k	2.157 k-m
	100 mg ^l ⁻¹	Sakha-93	12.33 f-i	16.91 fg	14.62 h	2.300 h-l	3.600 ef	2.950 d-f
		Gemiza-7	16.33 bc	21.42 b	18.88 bc	3.703 c	4.053 cd	3.878 c
		Gemiza-9	13.32 e-g	17.05 fg	15.19 f-h	2.390 h-j	3.797 de	3.093 d
Thiamine	50 mg ^l ⁻¹	Sakha-93	15.83 bc	16.27 gh	16.05 ef	2.200 i-m	3.783 de	2.992 de
		Gemiza-7	11.72 g-k	13.76 kl	12.74 i-k	1.920 k-o	3.043 g-k	2.482 h-j
		Gemiza-9	15.56 cd	15.24 ij	15.40 f-h	1.953 j-n	3.293 f-h	2.623 g-i
	100 mg ^l ⁻¹	Sakha-93	11.38 h-k	13.87 kl	12.62 i-k	2.187 i-m	3.033 g-k	2.610 g-i
		Gemiza-7	9.453 l-o	11.51 op	10.48 l	1.830 m-o	2.680 k	2.255 j-l
		Gemiza-9	12.17 f-i	14.54 jk	13.35 i	2.753 e-h	3.080 g-k	2.917 d-g
Ascorbic Acid	100 mg ^l ⁻¹	Sakha-93	10.88 i-l	13.83 kl	12.36 i-k	2.677 f-h	3.157 f-j	2.917 d-g
		Gemiza-7	13.06 f-h	17.56 ef	15.31 f-h	3.850 bc	4.087 cd	3.968 c
		Gemiza-9	9.387 l-o	11.95 no	10.67 l	2.070 j-n	2.193 l	2.132 k-m
	100 mg ^l ⁻¹	Sakha-93	12.15 f-j	18.56 d	15.36 f-h	2.730 f-h	3.260 f-h	2.995 de
		Gemiza-7	19.36 a	22.75 a	21.06 a	4.720 a	5.127 a	4.923 a
		Gemiza-9	17.64 b	18.64 d	18.14 cd	3.803 bc	4.210 b-d	4.007 c
Yeast	1000 mg ^l ⁻¹	Sakha-93	10.60 i-m	15.03 j	12.81 ij	2.937 e-g	3.410 e-g	3.173 d
		Gemiza-7	14.94 c-e	18.30 de	16.62 e	4.187 b	4.547 b	4.367 b
		Gemiza-9	11.53 g-k	13.17 lm	12.35 i-k	2.547 g-i	2.883 h-k	2.715 e-h
	2000 mg ^l ⁻¹	Sakha-93	8.817 m-o	11.48 op	10.15 l	2.717 f-h	2.617 k	2.667 f-i
		Gemiza-7	12.03 g-j	13.49 lm	12.76 i-k	3.013 ef	2.917 h-k	2.965 d-f
		Gemiza-9	7.990 op	11.46 op	9.727 lm	1.770 m-o	2.917 i-k	2.265 j-l

S1 = first season, S2 = second season, Comb. = combined data.

Table 5. Spike length, number of spikes per plant and weight of spike of some wheat varieties, as affected by some growth bio-regulating substances during two growing seasons and their combined data.

Treatments		Spike length (cm)			No. of Tillers/plant			weight of spike (g)		
		S1	S2	Comb.	S1	S2	Comb.	S1	S2	Comb.
Control		7.15 c	6.87 e	7.01 e	1.75 e	2.33 d	2.044 e	1.84 e	2.77 e	2.30 h
Tryptophan	50 mg ^l ⁻¹	9.21 ab	12.14 a	10.68 a	3.92 a-c	5.56 a	4.74 a	4.39 a	6.42 a	5.40 a
	100 mg ^l ⁻¹	9.43 a	10.77 cd	10.10 bc	3.41 b-d	4.33 bc	3.87 b-d	3.81 b	5.42 cd	4.62 c-e
Cystine	100 mg ^l ⁻¹	8.55 b	11.33 bc	9.94 cd	2.82 d	3.78 c	3.30 d	3.36 cd	5.79 b-d	4.57 d-f
	150 mg ^l ⁻¹	9.04 ab	10.65 d	9.85 cd	3.17 cd	4.00 c	3.58 cd	3.13 d	5.34 d	4.23 g
Thiamine	50 mg ^l ⁻¹	9.26 ab	11.04 cd	10.15 bc	3.50 b-d	4.42 bc	3.96 b-d	3.11 d	5.63 b-d	4.37 e-g
	100 mg ^l ⁻¹	8.86 ab	12.19 a	10.52 ab	3.61 b-d	3.94 c	3.78 b-d	3.79 b	5.95 a-c	4.87 bc
Ascorbic Acid	50 mg ^l ⁻¹	8.89 ab	10.82 cd	9.85 cd	3.41 b-d	3.94 c	3.68 cd	3.80 b	6.12 ab	4.96 b
	100 mg ^l ⁻¹	8.68 ab	11.88 ab	10.28 a-c	4.04 ab	4.78 a-c	4.41 ab	3.23 d	5.66 b-d	4.44 d-g
Yeast	1000 mg ^l ⁻¹	8.77 ab	11.65 ab	10.21 a-c	4.44 a	5.22 ab	4.83 a	3.52 c	5.90 a-c	4.71 b-d
	1500 mg ^l ⁻¹	8.51 ab	10.62 d	9.56 d	3.72 a-c	4.50 bc	4.11 bc	3.33 cd	5.24 d	4.29 fg
Sakha-93		8.09 c	9.96 c	9.023 c	3.56 a	4.33 ab	3.94 a	2.75 c	5.02 c	3.88 c
Gemiza-7		9.81 a	12.31 a	11.06 a	3.30 a	3.98 b	3.64 b	3.84 a	6.09 a	4.96 a
Gemiza-9		8.38 b	10.45 b	9.416 b	3.46 a	4.45 a	3.95 a	3.59 b	5.32 b	4.46 b

S1 = first season, S2 = second season, Comb. = combined data.

highest values of number of spikes per plant in the second season and the combined data, number of grains per spike in the first season and the combined data and straw yield in the two growing season and their combined data.

Whereas, Sakha-93 variety was exceeded in number of spikes per plant in the first season.

Effect of interaction

The effect of interaction between foliar application of bio-regulating

substances and the three wheat varieties on yield and its components was significant and highly increased all yield components compared with the untreated plants (Tables 7-8).

Data indicated that, the highest

Table 6. Number of grains per spike, grainsweight per spike, grain and straw yield per feddanof some wheat varieties, as affected by some growth bio-regulating substances during two growing seasons and their combined data.

Treatments	No. of grains/spike			Grains weight(g)/spike			Grain yield (Tons/feddan)			Straw yield (Tons/feddan)			
	S1	S2	Comb.	S1	S2	Comb.	S1	S2	Comb.	S1	S2	Comb.	
Control	30.01 g	29.63 d	29.82 e	1.33 f	1.94 e	1.64 g	1.61 f	2.00 f	1.80 f	1.82 g	2.37 f	2.09 h	
Tryptophan	50 mg/l-1	55.48 a	69.44 ab	62.46 a	2.85 a	5.18 a	4.01 a	2.59 a	3.02 a	2.80 a	4.26 bc	4.95 b	4.60 b
	100 mg/l-1	44.93 d	59.47 c	52.20 cd	2.44 bc	4.14 d	3.29 c-e	2.37 bc	2.73 b	2.55 b	3.49 e	4.02 d	3.75 e
Cystine	100 mg/l-1	45.47 cd	72.14 a	58.80 b	2.22 cd	4.81 a-c	3.51 bc	1.83 e	2.30 e	2.06 e	2.88 f	3.61 e	3.25 g
	150 mg/l-1	38.61 f	60.08 c	49.34 d	1.83 e	4.14 d	2.98 f	2.33 bc	2.46de	2.40 cd	4.19 c	4.43 c	4.31 c
Thiamine	50 mg/l-1	41.07 e	65.04 bc	53.05 c	2.01 de	4.56 b-d	3.29 c-e	2.17 d	2.50 c-e	2.34 d	3.76 de	4.29 cd	4.02 d
	100 mg/l-1	50.44 b	67.73 ab	59.08 b	2.61 ab	4.91 ab	3.76 ab	2.32 c	2.63b-d	2.48 bc	4.53 b	5.04 b	4.79 b
Ascorbic Acid	50 mg/l-1	47.81 c	70.35 ab	59.08 b	2.42 bc	4.96 ab	3.69 b	2.11 d	2.47 de	2.29 d	4.00 cd	4.51 c	4.25 c
	100 mg/l-1	40.34 ef	64.92 bc	52.63 c	2.09 c-e	4.36 cd	3.23 d-f	2.42 b	2.69 bc	2.56 b	4.97 a	5.63 a	5.30 a
Yeast	1000 mg/l-1	47.14 cd	65.32 bc	56.23 b	2.20 cd	4.82 a-c	3.51 b-d	2.32 c	3.14 a	2.73 a	2.99 f	4.05 d	3.52 f
	1500 mg/l-1	41.28 e	60.69 c	50.99 cd	1.90 de	4.20 d	3.05 ef	2.09 d	2.52 b-d	2.31 d	2.80 f	3.43 e	3.12 g
Sakha-93		42.97 b	61.51 a	52.24 b	1.84 b	3.89 c	2.86 c	1.95 c	2.36 b	2.15 c	3.54 a	4.23 a	3.88 b
Gemiza-7		42.63 b	62.63 a	52.63 b	2.41 a	5.00 a	3.71 a	2.42 a	2.73 a	2.57 a	3.58 a	4.03 b	3.81 b
Gemiza-9		46.02 a	62.62 a	54.32 a	2.26 a	4.21 b	3.24 b	2.22 b	2.67 a	2.45 b	3.70 a	4.37 a	4.04 a

S1 = first season, S2 = second season, Comb. = combined data.

Table 7. Spike length, number of Spikes per plant and weight of spike of, as affected by interaction between some growth bio-regulating substances treatments and wheat varieties during two growing seasons and their combined data.

Treatments		Spike length (cm)			No. of Spikes/plant			weight of spike (g)			
		S1	S2	Comb.	S1	S2	Comb.	S1	S2	Comb.	
Control	Sakha-93	6.91 m	6.80 m	6.85 p	1.67 h	2.67 g-i	2.17 lm	1.69 o	2.79 l	2.24 n	
	Gemiza-7	7.41 j-m	7.07 m	7.24 p	1.93 gh	2.33 hi	2.13 lm	2.19 mn	2.68 l	2.44 n	
	Gemiza-9	7.14 lm	6.73 m	6.94 p	1.67 h	2.00 i	1.83 m	1.65 o	2.83 l	2.24 n	
Tryptophan	50 mg/l-1	Sakha-93	8.68 f-i	12.56 cd	10.62 e-g	3.25 b-f	4.50 b-f	3.87 c-k	3.78 d-g	6.78 ab	5.28 b-d
		Gemiza-7	10.24 ab	12.33 de	11.28 b-d	4.00 a-e	5.17 b-e	4.58 b-f	4.60 ab	6.29 b-f	5.28 b-d
		Gemiza-9	8.72 f-i	11.53 ef	10.13 f-i	4.50 ab	7.00 a	5.75 a	4.78 a	6.19 b-g	5.48 ab
	100 mg/l-1	Sakha-93	8.93 e-g	9.67 i-l	9.30 k-n	3.33 b-f	3.50 f-h	3.42 i-k	3.25 hi	5.30 hi	4.27 ij
		Gemiza-7	10.16 bc	11.34 f	10.75 d-f	3.06 c-g	4.00 c-g	3.53 g-k	4.12 c-e	5.50 f-i	4.81 d-h
		Gemiza-9	9.21 c-f	11.30 f	10.26 f-h	3.83 a-e	5.50 bc	4.67 b-e	4.07 c-e	5.48 f-i	4.77 e-h
Cysteine	50 mg/l-1	Sakha-93	7.97 g-l	10.57 f-j	9.27 k-n	3.25 b-f	4.00 c-g	3.62 f-k	3.45 g-i	5.04 ij	4.24 ij
		Gemiza-7	9.84 b-e	12.82 b-d	11.33 b-d	2.33 f-h	3.50 f-h	2.92 kl	3.44 g-i	6.46 a-d	4.95 c-f
		Gemiza-9	7.84 h-m	10.61 f-i	9.23 k-n	2.89 e-g	3.83 d-g	3.36 i-k	3.18 ij	5.88 c-i	4.53 f-i
	100 mg/l-1	Sakha-93	8.62 f-i	9.567 kl	9.09 l-n	3.33 b-f	4.17 c-g	3.75 d-k	2.58 k-m	4.46 jk	3.52 lm
		Gemiza-7	10.19 bc	12.77 cd	11.48 bc	3.17 c-g	4.00 c-g	3.58 g-k	3.61 e-i	6.28 b-f	4.95 c-f
		Gemiza-9	8.32 f-k	9.60 j-l	8.96mn	3.00 d-g	3.83 d-g	3.42 i-k	3.21 ij	5.27 hj	4.24 ij
Thiamine	50 mg/l-1	Sakha-93	8.85 e-h	9.66 i-l	9.25 k-n	4.00 a-e	5.00 b-f	4.50 b-h	2.27 l-n	5.07 ij	3.67 kl
		Gemiza-7	11.14 a	12.53 d	11.84 ab	3.50 b-f	4.27 c-f	3.88 c-k	3.54 f-i	6.57 a-c	5.06 b-e
		Gemiza-9	7.80 i-m	10.93 f-h	9.36 k-n	3.00 d-g	4.00 c-g	3.50 i-k	3.51 f-i	5.26 hi	4.39 h-j
	100 mg/l-1	Sakha-93	8.37 f-j	10.58 f-j	9.47 i-m	3.50 b-f	3.83 d-g	3.67 f-k	3.63 e-i	5.29 hi	4.46 f-j
		Gemiza-7	9.95 b-d	14.87 a	12.41 a	4.17 a-e	4.33 b-f	4.25 c-j	3.90 c-g	6.84 ab	5.37 a-c
		Gemiza-9	8.26 f-k	11.12 fg	9.69 h-l	3.17 c-g	3.67 e-h	3.42 i-k	3.84 d-g	5.72 d-i	4.78 e-h
Ascorbic Acid	100 mg/l-1	Sakha-93	7.43 j-m	10.06 h-k	8.74 n	3.42 b-f	4.00 c-g	3.71 e-k	2.76 jk	5.62 e-i	4.19 ij
		Gemiza-7	10.40 ab	11.57 ef	10.98 c-e	3.11 c-g	3.50 f-h	3.30 jk	4.38 a-c	7.12 a	5.75 a
		Gemiza-9	8.84 e-h	10.82 f-h	9.83 j-n	3.71 b-e	4.33 b-f	4.02 c-j	4.26 b-d	5.62 e-i	4.94 c-f
	100 mg/l-1	Sakha-93	7.70 i-m	11.14 fg	9.42 j-n	4.03 a-e	5.00 b-f	4.52 b-g	2.70 kl	5.39 g-i	4.04 jk
		Gemiza-7	9.72 b-e	13.49 bc	11.60 bc	3.83 a-e	4.17 c-g	4.00 c-j	3.75 e-h	6.02 b-h	4.88 d-g
		Gemiza-9	8.61 f-i	11.01 f-h	9.81 h-k	4.25 a-d	5.17 b-e	4.71 b-d	3.24 i	5.07 e-i	4.41 g-j
Yeast	1000 mg/l-1	Sakha-93	8.21 f-k	10.16 g-k	9.18 k-n	5.00 a	5.83 ab	5.42 ab	1.90 no	5.34 hi	3.62 kl
		Gemiza-7	9.03 d-f	13.73 b	11.38 b-d	4.17 a-e	4.50 b-f	4.33 c-i	4.67 ab	6.81 ab	5.74 a
		Gemiza-9	9.06 d-f	11.06 fg	10.06 g-j	4.17 a-e	5.33 b-d	4.75 bc	3.98 c-f	5.53 f-i	4.76 e-h
	2000 mg/l-1	Sakha-93	7.32 k-m	8.75 l	8.03 o	4.33 a-c	5.17 b-e	4.75 bc	2.19 mn	4.16 k	3.17 m
		Gemiza-7	9.83 b-e	12.84 b-d	11.34 b-d	3.00 d-g	4.00 c-g	3.50 h-k	4.01 c-f	6.38 a-e	5.19 b-e
		Gemiza-9	8.37 f-j	10.26 g-k	9.32 k-n	3.83 a-e	4.33 b-fj	4.08 c-j	3.79 d-g	5.19 h-j	4.49 f-j

S1 = first season, S2 = second season, Comb. = combined data.

value of spike length was obtained by thiamine treatment when used at 50 mg^l⁻¹ concentration in the first season and 100 mg^l⁻¹ in the second season and combined data when used with Gemiza-7 variety. The highest value of number of spikes per plant in the first season had been resulted by the interaction between 1000 mg^l⁻¹ yeast extract with Sakha-93 variety, while it had been resulted from interaction between 50 mg^l⁻¹ of tryptophan with

Gemiza-9 variety during the second season and the combined data analysis.

The most effective treatment in the mean values of spike weight was that of 50 mg^l⁻¹ tryptophan with Gemiza-9 variety in the first season. On the other side ascorbic acid treatment at 50 mg^l⁻¹ was more effective on the weight of spike when used on Gemiza-7 during the second season and the combined data analysis. Results showed that, the most effective treatment on grains weight per

spike in the first season was that of 50 mg^l⁻¹ tryptophan on Gemiza-7 variety, while, ascorbic acid was more effective in the second season and the combined data when used at 50 mg^l⁻¹ concentration on the same variety.

Foliar application of 50 mg^l⁻¹ tryptophan on Gemiza-9 variety recorded the highest values in grain yield in the first, second season and their combined data.

The best treatments of straw yield

Table 8. Number of grains per spike, grains weight per spike and grain and straw yield as affected by interaction between some growth bio-regulators treatments and varieties of wheat, during two growing seasons and their combined data.

Treatments	No. of grains/spike			Grains weight(g)/spike			Grain yield (Tons/feddan)			Straw yield (Tons/feddan)			
	S1	S2	Comb.	S1	S2	Comb.	S1	S2	Comb.	S1	S2	Comb.	
	Control	Sakha-93 37.63 l Gemiza-7 27.00 n Gemiza-9 25.40 n	32.67 h 25.47 h 30.75 h	35.15 m 26.23 n 28.08 n	1.13 o 1.29 no 1.58 k-o	2.28 m 1.72 m 1.82 m	1.71 n 1.50 n 1.70 n	1.55 o 1.72 l-o 1.55 o	1.88 m 2.06 lm 2.07 k-m	1.71 q 1.89 o-q 1.81 q	1.50 m 1.75 lm 2.22 k-m	2.04 r 2.09 r 2.97 q	1.77 p 1.92 p 2.59 o
Tryptophan	Sakha-93 54.12 bc Gemiza-7 53.00 b-d Gemiza-9 59.33 a Sakha-93 41.73 i-k Gemiza-7 46.53 e-h Gemiza-9 46.53 e-h	73.83 a-d 58.17 fg 76.33 a 59.67 e-g 57.83 fg 60.92 e-g	63.97 ab 55.58 c-i 67.83 a 50.70 i-k 52.18 g-j 53.72 d-j	2.69 a-e 3.08 a 2.78 a-c 1.97 g-m 2.67 a-f 2.67 a-f	5.80 ab 4.70 d-i 5.04 b-f 3.83 j-l 4.43 e-j 4.17 g-j	4.24 ab 3.89 b-f 3.91 b-f 2.90 j-l 3.55 d-h 3.42 f-i	2.49 d-i 2.78 bc 3.13 a 2.19 k-n 2.42 g-j 3.03 a	2.35 g 2.66 a-c 2.75 a 2.08 hi 2.35 fg 2.67 ab	2.64 e-i 2.91 c-f 3.51 a 2.30 h-l 2.48 f-k 3.40 ab	2.49 d-i 2.78 bc 3.13 a 2.19 k-n 2.42 g-j 3.03 a	4.46 a-d 4.17 b-e 4.15 b-e 3.29 e-i 3.39 e-i 3.78 d-g	5.00 a-f 4.56 d-j 5.28 a-d 3.63 l-q 3.59 m-q 4.83 c-h	4.73 c-f 4.36 e-i 4.71 c-g 3.46 k-m 3.49 k-m 4.31 e-j
Cysteine	Sakha-93 48.20 ef Gemiza-7 46.53 e-h Gemiza-9 41.67 i-k Sakha-93 38.53 kl Gemiza-7 39.50 j-l Gemiza-9 37.80 l	71.23 a-e 75.19 a-c 70.00 a-f 53.08 g 67.58 a-f 59.57 e-g	59.72 b-e 60.86 bc 55.83 c-i 45.81 kl 53.54 e-j 48.68 l-j	1.81 i-n 2.51 a-g 2.35 b-i 1.64 j-o 2.08 f-l 1.76 l-n	3.85 i-l 5.60 a-c 4.97 b-g 3.32 kl 5.14 b-e 3.97 h-k	2.83 j-l 4.06 b-d 3.66 d-g 2.48 lm 3.61 d-h 2.87 j-l	1.87 pq 2.25 j-n 2.07 m-p 2.08 m-p 2.63 c-g 2.47 e-j	1.68 m-o 1.93 h-k 1.88 i-l 1.97 h-j 2.61 a-d 2.42 d-g	2.06 lm 2.57 f-j 2.27 i-m 2.19 j-m 2.66 e-i 2.53 f-j	1.87 pq 2.25 j-n 2.07 m-p 2.08 m-p 2.63 c-g 2.47 e-j	3.16 f-j 2.77 h-k 2.71 h-k 4.25 b-e 4.12 b-e 4.19 b-e	3.89 j-p 3.68 k-q 3.27 o-q 4.70 d-i 4.20 g-n 4.38 e-l	3.52 k-m 3.23 l-n 2.99 m-o 4.47 d-h 4.16 f-j 4.29 e-j
Thiamine	Sakha-93 47.27 e-g Gemiza-7 37.13 l Gemiza-9 38.80 kl Sakha-93 53.00 b-d Gemiza-7 48.33 ef Gemiza-9 50.00 de	62.42 c-g 65.39 a-g 67.31 a-f 67.50 a-f 67.11 a-f 68.57 a-f	54.84 c-j 51.26 h-k 53.05 f-j 60.25 b-d 57.72 b-h 59.28 b-f	1.40 m-o 2.33 b-i 2.30 b-i 2.53 a-g 2.74 a-d 2.55 a-g	3.94 h-l 5.55 a-c 4.20 f-j 4.34 e-j 5.61 a-c 4.79 c-h	2.67 k-m 3.94 b-e 3.25 g-j 3.43 e-i 4.17 a-c 3.67 d-g	2.19 k-n 2.72 cd 2.10 l-o 2.45 f-j 2.80 bc 2.17 k-n	2.13 h 2.44 d-g 1.95 h-k 2.36 fg 2.60 a-e 2.00 h-j	2.24 i-m 2.99 c-e 2.26 i-m 2.55 f-j 3.00 c-e 2.34 g-l	2.19 k-n 2.72 cd 2.10 l-o 2.45 f-j 2.80 bc 2.17 k-n	4.36 e-l 4.09 h-n 4.40 e-k 5.02 a-f 5.00 a-f 5.11 a-e	4.25 e-j 3.72 j-l 4.10 h-j 4.61 c-h 4.79 b-e 4.81 a-c	4.25 e-j 3.72 j-l 4.10 h-j 4.61 c-h 4.79 b-e 4.96 a-d
Ascorbic Acid	Sakha-93 41.69 i-k Gemiza-7 45.82 f-h Gemiza-9 55.93 b Sakha-93 31.63 m Gemiza-7 42.97 h-j Gemiza-9 46.42 e-h	70.08 a-f 75.58 ab 65.39 a-g 69.58 a-f 62.08 d-g 63.08b-g	55.89 c-i 60.70 bc 60.66 bc 50.61 i-k 52.53 g-j 54.75 c-j	1.89 h-m 2.88 ab 2.48 a-h 1.84 i-n 2.24 c-j 2.21 c-j	4.16 g-j 6.20 a 4.53 d-j 3.92 i-l 4.98 b-g 4.19 f-j	3.02 i-k 4.54 a 3.50 e-i 2.88 j-l 3.61 d-h 3.20 g-j	2.03 n-p 2.54 d-h 2.30 i-m 2.33 h-l 2.66 c-f 2.69 c-e	1.82 j-m 2.40 e-g 2.11 h 2.12 h 2.55 b-f 2.61 a-d	2.24 i-m 2.67 e-i 2.49 f-k 2.54 f-j 2.78 d-f 2.77 d-g	2.03 n-p 2.54 d-h 2.30 i-m 2.33 h-l 2.66 c-f 2.69 c-e	4.27 f-n 4.95 b-g 4.31 f-m 5.65 ab 5.51 a-c 5.73 a	4.27 f-n 4.95 b-g 4.31 f-m 5.65 ab 5.51 a-c 5.31 ab	3.87 i-k 4.78 b-e 4.12 g-j 5.45 a 5.14 a-c 5.31 ab
Yeast	Sakha-93 45.20 f-i Gemiza-7 43.80 g-i Gemiza-9 52.43 b-d Sakha-93 33.63 m Gemiza-7 38.33 kl Gemiza-9 51.87 cd	62.28 d-g 68.90 a-f 64.78 a-g 54.25 g 65.67 a-g 62.17 d-g	53.74 d-j 56.35 c-i 58.60 b-g 43.94 l 52.00 g-j 57.02 c-i	1.81 l-n 2.61 a-f 2.16 d-k 1.52 l-o 2.10 e-l 2.08 e-l	4.21 f-j 5.75 ab 4.49 d-j 3.13 l 5.32 b-d 4.15 g-j	3.01 i-k 4.18 a-c 3.32 g-j 2.32 m 3.71 c-g 3.11 h-k	2.45 f-j 2.97 ab 2.76 bc 1.91 o-q 2.66 c-f 2.36 h-k	2.40 e-g 2.73 ab 2.47 c-g 1.62 no 2.59 a-e 2.07 hi	2.67 e-i 3.22 a-c 3.06 b-e 2.20 j-m 2.72 e-h 2.65 e-i	2.45 f-j 2.97 ab 2.76 bc 1.91 o-q 2.66 c-f 2.36 h-k	4.01 i-n 3.52 n-q 4.60 d-j 3.98 i-o 3.10 q 3.21 pq	4.01 i-n 3.52 n-q 4.60 d-j 3.98 i-o 3.10 q 3.21 pq	3.13 l-o 3.25 l-n 4.16 f-j 3.46 k-m 3.02 m-o 2.86 no

S1 = first season, S2 = second season, Comb. = combined data.

was foliar application of 100 mg l⁻¹ ascorbic acid on Sakha-93 variety during the first season and combined data and the same treatment with Gemiza-9 in the second season.

The total photosynthetic pigments

Table 9. Average values of the photosynthetic pigments content (mg/g fresh weight of wheat leaves) , as affected by some growth bio-regulators treatments and varieties of wheat. (Average of two seasons).

Treatments		Chlorophyll (a) mg/g	Chlorophyll (b) mg/g	Carotenoids mg/g
Control		0.6661 j	0.3028 h	0.2219 h
Tryptophan	50 mg l ⁻¹	1.229 a	0.6101 a	0.4390 a
	100 mg l ⁻¹	1.021 e	0.5308 c	0.3901 bc
Cystine	100 mg l ⁻¹	0.6857 i	0.3446 g	0.2723 g
	150 mg l ⁻¹	0.8428 h	0.4350 f	0.3169 ef
Thiamine	50 mg l ⁻¹	0.8422 h	0.4370 f	0.3092 f
	100 mg l ⁻¹	1.173 b	0.5323 c	0.3977 b
Ascorbic Acid	50 mg l ⁻¹	0.8592 g	0.5052 d	0.3218 e
	100 mg l ⁻¹	1.092 c	0.5607 b	0.3847 c
Yeast	1000 mg l ⁻¹	1.037 d	0.4882 e	0.4434 a
	1500 mg l ⁻¹	0.8741 f	0.4360 f	0.3649 d
Sakha-93		0.863 b	0.4673 b	0.3244 b
Gemiza-7		0.971 a	0.4746 a	0.3660 a
Gemiza-9		0.981 a	0.4715 ab	0.3628 a

S1 = first season, S2 = second season, Comb. = combined data.

Table 10. Effect of interaction between wheat varieties and bio-regulator treatments on the photosynthetic pigments content (mg/g fresh weight of wheat leaves) (average of two successive seasons).

Treatments		Chlorophyll (a) mg/g	Chlorophyll (b) mg/g	Carotenoids mg/g	
Control	Sakha-93	0.540 n	0.292 o	0.213 q	
	Gemiza-7	0.626 m	0.294 o	0.211 q	
	Gemiza-9	0.832 jk	0.323 n	0.242 op	
Tryptophan	50 mg l ⁻¹	Sakha-93	1.263 b	0.638 a	0.437 de
		Gemiza-7	1.261 b	0.624 a	0.479 b
		Gemiza-9	1.162 cd	0.568 cd	0.401 g
	100 mg l ⁻¹	Sakha-93	0.993 f	0.526 e	0.365 ij
		Gemiza-7	1.168 c	0.555 d	0.426 ef
		Gemiza-9	0.900 g-i	0.511 e	0.379 hi
Cysteine	50 mg l ⁻¹	Sakha-93	0.544 n	0.344 m	0.203 q
		Gemiza-7	0.821 jk	0.379 l	0.370 i
		Gemiza-9	0.692 l	0.311 n	0.244 op
	100 mg l ⁻¹	Sakha-93	0.821 jk	0.433 j	0.287 n
		Gemiza-7	0.932 gh	0.435 j	0.382 hi
		Gemiza-9	0.776 k	0.437 j	0.281 n
Thiamine	50 mg l ⁻¹	Sakha-93	0.869 ij	0.464 gh	0.319 m
		Gemiza-7	0.939 fg	0.464 gh	0.350 jk
		Gemiza-9	0.718 l	0.384 l	0.259 o
	100 mg l ⁻¹	Sakha-93	0.910 g-i	0.492 f	0.324 m
		Gemiza-7	1.473 a	0.526 e	0.456 c
		Gemiza-9	1.135 cd	0.579 c	0.413 fg
Ascorbic Acid	100 mg l ⁻¹	Sakha-93	0.778 k	0.481 fg	0.278 n
		Gemiza-7	0.638 m	0.479 fg	0.237 p
		Gemiza-9	11.161 cd	0.556 d	0.451 cd
	100 mg l ⁻¹	Sakha-93	0.874 h-j	0.515 e	0.317 m
		Gemiza-7	0.959 fg	0.564 cd	0.342 kl
		Gemiza-9	1.443 a	0.603 b	0.495 ab
Yeast	1000 mg l ⁻¹	Sakha-93	1.074 e	0.493 f	0.501 a
		Gemiza-7	0.930 gh	0.458 hi	0.401 g
		Gemiza-9	1.108 de	0.513 e	0.429 ef
	2000 mg l ⁻¹	Sakha-93	0.828 jk	0.465 gh	0.326 lm
		Gemiza-7	0.927 gh	0.443 ij	0.373 i
		Gemiza-9	0.867 ij	0.400 k	0.396 gh

S1 = first season, S2 = second season, Comb. = combined data.

Effect of bio-regulator treatments

Data presented in Table (9) showed that, total photosynthetic pigments (chlorophyll a, b and carotenoids) were significantly increased as affected by bio-regulating substances compared to the

untreated plants (control) at the two growth stages.

Tryptophan was more effective in all studied characters, when used at the concentration of 50 mg l⁻¹ except the total carotenoids which was more affected by 1000 mg l⁻¹ of yeast extract.

Varieties performance

There were no significant differences between Gemiza-7 and Gemiza-9 varieties in chlorophyll (a) content. The highest value was obtained by Gemiza-9 variety during heading stage. While, the last variety was Sakha-93 which recorded the lowest value in this parameter. Gemiza-7 variety exceeded the other two varieties in chlorophyll (b) during heading stage.

Concerning total carotenoids in the leaves Gemiza-7 variety exceeded the other two varieties in this respect during heading stage.

Effect of interaction

Data in Table (10) indicated that, thiamine treatment when used at 100 mg l⁻¹ concentration with Gemiza-7 variety resulted in the highest value of chlorophyll (a) at the heading stage.

The highest value of chlorophyll (b) was produced from interaction between 50 mg l⁻¹ tryptophan with Sakha-93 variety during heading stage.

As for the effect of the interaction on the carotenoids contents in the leaves of wheat plants, the most effective treatments was 1000 mg l⁻¹ of yeast extract with Sakha-93 variety at heading stage.

4 Discussion

Effect of bio-regulators on growth

According to the present data, the effect of foliar application of the different applied bioregulating substances at different concentrations on plant height, number of tillers and leaves per plant and fresh and dry weight of plant at heading stage were high significantly increased in both season and their combined data of all bio-regulator treatments. These results are in harmony with those obtained by Gamal El-Din and Zaki (2005) who found that, at harvest stage all treatments of (Lysine, phenylalanine

and L-cysteine) led to significant increases in plant height, number of pods and seeds per plant, dry weight of pods of lupine plants. EL-Bassiouny (2005) found that application of nicotinamide or tryptophan resulted in significant increases in plant growth and grain yield, concomitantly with an increase in the level of IAA, GA₃, cytokinins, photosynthetic pigments and a decrease in ABA content of wheat plant.

On the other hand, Gadalla, (2009) reported that, application of ascorbic acid increased significantly flag leaf area in wheat plants grown under salinity conditions. These results may be due to the effect of ascorbic acid on osmotic adjustment and maintaining leaf turgor potential as a consequence of increasing leaf water potential and relative water content as compared to control plants. Farouk, (2011).

Yield and yield components

As for the effect of bio-regulating substances on yield and yield components the present data show that, foliar application of all used concentrations of bio-regulating substances significantly increased the yield and yield components of all used wheat varieties. Similarly, El-Bassiouny, (2005) reported that application of tryptophan resulted in significant increase in plant growth and grain yield, concomitantly with an increase in the level of IAA, GA₃, cytokinins, photosynthetic pigments and a decrease in ABA content of wheat plant. Also, Abdel-Halim (1995) reported that, ascorbic acid at 100 and 200 ppm led to increased contents of indols (IAA, IAN and IBA) in the shoot of tomato. Amin et al., (2008) reported that ascorbic acid recorded the highest increase in straw yield per plant and per feddan of wheat plant cv. Gemiza10.

These results might emphasize the role of ascorbic acid as scavenge the free radicals which caused increase in the oxidation in plant tissues (Noctor and Foyer, 1998).

Photosynthetic pigments

It is observed from the presented data that, total photosynthetic pigments (chlorophyll a, b and carotenoids) were significantly increased as affected by

the different bio-regulating substances compared to untreated plants (control), these results are true for all photosynthetic pigments.

The induced effect of tryptophan on chlorophyll biosynthesis may be due to its role in IAA biosynthesis (Arshad et al., 1995; Barazani and Friedman, 2000). Also, the positive effect of foliar application of amino acids in enhancing all photosynthetic pigments percentage might be attributed to its positive effect on the succinyl COA (Kerb's cycle intermediate) and their effect in initiating the biosynthetic pathway leading to chlorophyll formation.

References

1. Abdel-Halim SM. Effect of some vitamins as growth regulators on growth, yield and endogenous hormones of tomato plants during winter. *Egypt J Appl Sci* (1995) 10(2): 322-334.
2. Amer SSA. Growth, green pods yield and seeds yield of common bean (*Phaseolus vulgaris* L) as affected by active dry yeast, salicylic acid and their interaction. *J Agric Sci, Mansoura Univ* (2004) 29(3): 1407-1422.
3. Amin AA, M El-Sh Rashad, FAE Gharib. Changes in morphological, physiological and reproductive characters of wheat plants as affected by foliar application with salicylic acid and ascorbic acid. *Australian J Basic Appl Sci* (2008) 2(2): 252-261.
4. Arshad M, A Hussain, A Shakoor. Effect of soil applied L-tryptophan on growth and chemical composition of cotton. *J Plant Nutr* (1995) 18: 317-329.
5. Barazani O, J Friedman. Effect of exogenously applied L-tryptophan on allelochemical activity of plant – growth promotion rhizobacteria. *J Chem Ecol* (2000) 26: 343-349.
6. Bidwell RGS. *Plant Physiology* (2nd Ed). MacMillan Publishing Co Inc, New York (1979): 236-238
7. Buchanan BB, W Gruissem, R Jones. *Biochemistry and Molecular Biology of Plants*. American Society of Plant Physiologists, Rockville, MD (2000).
8. Cakmak I, H Ozkan, HJ Braun, RM Welch, V Romheld. Zinc and iron concentrations in seeds of wild, primitive, and modern wheats. *Food Nutr Bull* (2000) 21: 401-403.
9. Duncan DB. Multiple Range and Multiple F-test. *Biometrics* (1955) 11: 1-42.
10. EL-Bassiouny HMS. Physiological Responses of Wheat to Salinity Alleviation by Nicotinamide and Tryptophan. *Int J Agri Biol* (2005) 7(4): 653-659.
11. Farouk S. Osmotic adjustment in wheat flag leaf in relation to flag leaf area and grain yield per plant. *J Stress physiol Biochem* (2011) 7(2): 117-138.
12. Gadalla SF. The role of antioxidants in inducing wheat flag leaf osmotic adjustment under salinity stress. *J Agric Sci Mansoura Univ* (2009) 34 (11): 10663 – 10685.
13. Gamal El-Din KM, NM Zaki. Effect of some amino acids on growth and biochemical constituents of lupine plant. (*Lupinus termis* L) *Egypt J Appl Sci* (2005) 20(6A): 70-78.
14. Huang D, W Wu, SR Abrams, AJ Cutler. The relationship of drought related geneexpression in *Arabidopsis thaliana* to hormonal and environmental factors. *J Exp Bot* (2008) 59: 2991-2997.
15. Kawasaki T. *Modern Chromatographic Analysis of Vitamins* (2nd Ed). New York, NY: Marcel Dekker, Inc (1992) 60: 319-354.
16. Kurtzman CP, JW Fell. *Biodiversity and Ecophysiology of Yeasts* (In: *The Yeast Handbook*, Gabor P, de la Rosa CL, eds) Berlin, Springer (2005): 11-30.
17. MacRitchie F. Physicochemical properties of wheat proteins in relation to functionality. *Adv Food Nutr Res* (1994) 36: 1-87.
18. Morgan PW. Effects of abiotic stresses on plant hormone systems. In *Stress Responses in Plants, Adaptation and Acclimation Mechanisms* (RC Alscher, JR Cumming, Eds), Wiley-Liss Inc, NY (1990): 113-146.
19. MSTATC. A Microcomputer Program for the Design Management and Analysis of Agronomic Research Experiments. Michigan State Univ USA. (1989)
20. Noctor G, CH Foyer. Ascorbate and glutathione: Keeping active oxygen under control. *Ann Rev Plant Physiol Plant Mol Biol* (1998) 49: 249-279.
21. Revenga C, J Brunner, N Henninger, K Kassem, R Payne. Pilot analysis of global ecosystems: Freshwater ecosystems. *W Resour Inst, Washington, DC* (2000): 83.
22. Sharaan AN, FS Abd El-Samie, IA Abd El-Gawad. Response of wheat varieties (*Triticum aestivum* L) to some environmental influences II: Effect of planting date and drought at different plant stages on yield and its components. *Porc Ninth Conf Agron Minufia Univ* (2000): 1-15.
23. Snedecor GW, WG Cochran. *Statistical Methods* (Seventh Edition). Ames, IA: Iowa State University Press (1980).
24. Sultan MS, AN Attia, AM Salma, SA El-Moursy, EM Said, MM Abou El-Nagah. Response of some wheat cultivars to planting and harvesting dates under different seed rates. *Proc 9th Conf Agron Minufiya Univ, 2-3 Sept of Abst* (2000): 15.
25. Wettstein D. Chlorophyll lethal und der submikro-skopicheformwechsel der plastiden. *Expt Cell Res* (1957) 12: 427-433.
26. Zaki NM, MA Ahmed, MS Hassanein. : Growth and yield of some wheat cultivars irrigated with saline water in newly cultivated land as affected by nitrogenfertilization. *Ann. Agric. Sci., Moshtohr* (2004) 42(2): 515-525.
27. Zaki NM, MS Hassanein, KM Gamal El-Din. Growth and yield of some wheatcultivars irrigated with saline water in newly cultivated land as affected by biofertilization. *J Appl Sci Res* (2007) 3(10): 1121-1126.